

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANT: Joseph M. Jilka
SERIAL NO: 10/086,062
FILED: February 28, 2002
TITLE: NOVEL PLANT PROMOTER SEQUENCES AND METHODS OF USE
FOR SAME

ART UNIT: 1635
EXAMINER: Epps, J.

131 DECLARATION OF JOSEPH M. JILKA

Commissioner of Patents and Trademarks
Washington, D.C. 20231

Dear Sir:

I, Joseph M. Jilka hereby declare the following:

1. That I am the inventor for the above-identified patent application; that I conceived and reduced to practice in the United States the invention claimed in the above-identified patent application prior to the international publication date of March 23, 2000, of the cited PCT Application No. WO 00/15810 to Goldsbrough as evidenced by the enclosed notebook pages.

2. Attached Exhibit A is a copy of notebook records relating to this conception wherein construction of proposed versions of the ubiquitin variants show a no heat shock version. Also relating to this conception is Exhibit B which is a copy of a table listing the promoters made which show a no heat shock version. Attached Exhibit C are primers among which is the no heat shock version, version 4A, 4B.

3. That pursuant to this conception, I actually reduced to practice in the United States the invention claimed in the above-identified patent application prior to March 23, 2000, the international publication date of the cited Goldsbrough patent. Attached Exhibit D and E are copies of the notebook records of Kathy Beifuss, who worked under my direction and supervision, however, did not contribute materially to the above-identified invention, relating to the actual reduction to practice, wherein Exhibit D shows use the no heat shock

EXHIBIT


DD

version in a mini-prep and Exhibit E shows use of the no heat shock version in sequencing. Additionally, attached Exhibits F and G relating to the actual reduction to practice is a copy of the notebook records of Chris Brooks and Elizabeth Wilfong, both who worked under my direction and supervision, however, did not contribute materially to the above-identified invention, showing the GUS reporter gene expression in corn seed using the Ubi promoter variant, GSC, the ubiquitin promoter having no heat shock elements. Wherein total soluble protein (1 μ g) was incubated in 100 μ l lysis buffer and the reaction initiated with 5mM 4-methylumbelliferyl β -D-glucuronide (MUG). The reaction was incubated for up to about 20 minutes at 37°C. At specific time points approximately 25 μ l of volume of the reaction mixture was transferred into a reading plate that had 175 μ l of Stop buffer in the well. The reaction plate was placed at 37°C until the next time point. Generally readings at 0, 15, 30, and 60 minutes were taken. Plates were read at 360nm excitation wavelength and 460 nm emission wavelength. GUS protein levels were then calculated by comparison to a standard curve of 1-100 μ M 4-methylumbelliferyl. Exhibit G shows results from a 10 minute reading. The dates of these records have been redacted, however, the acts of conception and reduction to practice occurred prior to March 23, 2000, the international publication date of the cited Goldsbrough patent.

4. That Exhibits, A, B, C, D, E, F, and G, which relate to the aforementioned conception and reduction to practice, correspond to the invention disclosed and claimed in the above-identified patent application.

5. The undersigned further declares that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application of any patent issuing thereon.

Date: 7/17/02


Joseph M. Jilka

Construction & utilization variants

Proposed versions

version 1.0 \longrightarrow A only
 version 2.0 \longrightarrow B only
 version 3.0 \longrightarrow A, B no overlap
 version 4.0 \longrightarrow no A, B

version

engas A8333C11

A8333C10

direct to 100 pmol / μ l

50 μ l each mixed

\downarrow
 25
 5
 40 slow cool

legation 2 μ l oligo (100 pmol)

2 μ l +23% aad

7 μ l H₂O

1.5 μ l 10X buffer


1.5 μ l 10mM ATP

10 μ l ligase

15 μ l

\downarrow 16° 1 hr

\downarrow 65° 15'



Joseph A. Fox
 Michael B. B.

ProtiGene Promoters

Name	Description	Reporter	Test vector	Transformation	Events
PGNpr1	maize polyubiquitin 1 (UBI1)	GUS	pPHIP8904	Corn	GSA
PGNpr2	maize globulin 1	GUS-Adh1s	pPGN7042	Corn	GSB
PGNpr3	maize 22 kD alpha-zelin	GUS-Adh1s	pPGN9075	Corn	
PGNpr4	maize UBI1 no heat shock elements (HSE): UbC	GUS-Adh1s	pPGN9071	Corn	PMD
PGNpr5	maize UBI1 no 3' HSE: UBI1D	GUS-Adh1s	pPGN7547	Corn	GSC
PGNpr6	maize UBI1 no 5' HSE: UBI1E	GUS-Adh1s	pPGN7565	Corn	GSD
PGNpr7	maize UBI1 no HSE overlap: Ubaf	GUS-Adh1s	pPGN7583	Corn	GSE
PGNpr8	maize UBI1 replace HSE with 3x P1 seed specific element: UbiG	GUS-Adh1s	pPGN7600	Corn	GSE
PGNpr9	teosinte polyubiquitin 1	GUS-Adh1s	pPGN8926	Corn	GSG
PGNpr10	teosinte polyubiquitin 1a	GUS-Adh1s	pPGN8984	Corn	GSI
PGNpr11	teosinte polyubiquitin 1	GUS-Adh1s	pPGN8985	Corn	GSM
PGNpr12	maize glutathione-S-transferase 1 (GST1)	GUS-Adh1s	pPGN8986	Corn	GSM
PGNpr13	synthetic promoter PsynD with 35S enhancer 5' (tested with maize Adh-1 Init)	GUS-Adh1s	pPGN9006	Corn	GSP
PGNpr14	synthetic promoter Psyn7 with 35S enhancer 5' (tested with maize Adh-1 Init)	GUS-Adh1s	pPGN9007	Corn	GSP
PGNpr15	maize HMG	GUS-Adh1s	pPGN9016	Corn	GSS
PGNpr16	maize P promoter (tested with maize Adh-1 Init)	GUS-Adh1s	pPGN9035	Corn	SCA
PGNpr17	modified version of Agrobacterium synthase (supMAS)	GUS	pPHIP10336	Corn	SCD
PGNpr18	bean pmsaath	GUS-10xGSP	pPGN9275	Pea	GSI
PGNpr19	maize UBI1 no 5' HSE with a little extended seq at 5' end (beyond p16)	GUS-Adh1s	pPGN5690	Pea	GSI
PGNpr20	rice glutelin 1 (3kb of 5' sequence)	GUS-Adh1s	pPGN9042	Corn	GSI
PGNpr21	rice glutelin 2 (3kb of 5' sequence)	GUS-Adh1s	pPGN9056	Corn	PMA
PGNpr22	rice globulin 26kDa	GUS-Adh1s	pPGN9057	Corn	PMB
PGNpr23	maize globulin 2	GUS-Adh1s	pPGN9060	Corn	PNC
PGNpr24	maize globulin 2	GUS-Adh1s	pPGN9076	Corn	



GIBCO BRL Custom Primers

Certificate of Analysis

Primer 1:

Primer Name: UBI HSP VER. 1A

Researcher:

Sequence (5' to 3'): PAG ACG GCA CGG CAT CTC TGT CGC TGC CTC CAC CGT TGG ACT TGC TOC GCT
GTC GGC ATC CAG AAA T

Primer Number: A8333C10 (C10)

Primer Length: 66

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 21299.2

Millimolar Extinction Coefficient: 678.6

Purity Desalted

Tm (1 M Na+) 96

Tm (50 mM Na+) 76

% GC 60

μg per OD: 31.3

nmoles per OD: 1.4

OD's 39.3

$\mu\text{g's}$ 1234

nmoles 67

Coupling Eff. 99%

Notes:

Primer 2:

Primer Name: UBI HSP VER.1B

Researcher:

Sequence (5' to 3'): PTT TCT GGA TGC CGA CAG CGG AGC AAG TCC AAC GGT GGA GGC AGC GAC AGA
GAT GCC GTG CCG TCT GC

Primer Number: A8333C11 (C11)

Primer Length: 67

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 21897.4

Millimolar Extinction Coefficient: 732.9

Purity Desalted

Tm (1 M Na+) 97

Tm (50 mM Na+) 78

% GC 62

μg per OD: 29.8

nmoles per OD: 1.3

OD's 10.7

$\mu\text{g's}$ 319

nmoles 14

Coupling Eff. 98%

Notes:


UBI 1A

57 nmoles

570 μl \rightarrow 100 $\mu\text{mol}/\mu\text{l}$

14 nmoles

140 μl \rightarrow 100 $\mu\text{mol}/\mu\text{l}$



* See Note about Quantities in Supporting Information.

LIFE TECHNOLOGIES.

GIBCO BRL Custom Primers

Certificate of Analysis

Primer 1:

Primer Name: UBI HSPA VER.2A

Primer Number: D0373B07 (B07)

Researcher:

Primer Length: 81

Sequence (5' to 3'): P-A GAC GGC ACG GCA TCT CTG TCG CTG CCT CTG GAC CCC TCT CGA CCA CCG
TTG GAC TTG CTC CGC TGT CGG CAT CCA GAA AT

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 26105.2

μg per OD: 31.6

Millimolar Extinction Coefficient: 824.3

nmoles per OD: 1.2

Purity	Desalt	OD's	90.0
Tm (1 M Na+)	98	$\mu\text{g's}^*$	2850
Tm (50 mM Na+)	77	nmoles	106
% GC	61	Coupling Eff.	98%

Notes:

10 μg \Rightarrow 100 pmol/ μmol

Primer 2:

Primer Name: UBI HSPB VER.2B

Primer Number: D0373B08 (B08)

Researcher:

Primer Length: 82

Sequence (5' to 3'): P-T TTC TGG ATG CCG ACA GCG GAG CAA GTC CAA CGG TGG TCG AGA GGG GTC
CAG AGG CAG CGA CAG AGA TGC CGT GCC GTC TGC

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 26872.4

μg per OD: 29.7

Millimolar Extinction Coefficient: 902.2

nmoles per OD: 1.1

Purity	Desalt	OD's	77.0
Tm (1 M Na+)	99	$\mu\text{g's}^*$	2294
Tm (50 mM Na+)	76	nmoles	85
% GC	63	Coupling Eff.	98%

Notes:

850 μg \Rightarrow 100 pmol/ μmol

Primer 3:

Primer Name: UBI HSPA VER.3A

Primer Number: D0373B09 (B09)

Researcher:

Primer Length: 81

Sequence (5' to 3'): P-A GAC GGC ACG GCA TCT CTG TCG CTG CCT CTC GAG AGT TCC GCT CCA CCG
TTG GAC TTG CTC CGC TGT CGG CAT CCA GAA AT

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 26160.2

μg per OD: 31.5

Millimolar Extinction Coefficient: 830.8


nmoles per OD: 1.2

Purity	Desalt	OD's	88.7
Tm (1 M Na+)	98	$\mu\text{g's}^*$	2783
Tm (50 mM Na+)	76	nmoles	106
% GC	60	Coupling Eff.	98%

Notes:

1040 μg \Rightarrow 100 pmol/ μmol

* See Note about Quantities in Supporting Information.



LIFE  TECHNOLOGIES.

GIBCO BRL Custom Primers Certificate of Analysis

Primer 4:

Primer Name: UBI HSPB VER.3B

Primer Number: D0373B10 (B10)

Researcher:

Primer Length: 82

Sequence (5' to 3'): P-T TTC TGG ATG CCG ACA GCG GAG CAA GTC CAA CGG TGG AGC GGA ACT CTC
GAG AGG CAG CGA CAG AGA TGC CGT GCC GTC TGC

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 26816.4

μg per OD: 29.7

Millimolar Extinction Coefficient: 901.3

nmoles per OD: 1.1

Purity	Desalt	OD's	83.2
Tm (1 M Na+)	99	$\mu\text{g's}$	2476
Tm (50 mM Na+)	77	nmoles	92
% GC	62	Coupling Eff.	98%
Notes:		930 μg \Rightarrow 100 μg	

Primer 5:

Primer Name: UBI HSPA VER.4A

Primer Number: D0373B11 (B11)

Researcher:

Primer Length: 86

Sequence (5' to 3'): P-A GAC GGC ACG GCA TCT CTG TCG CTG CCT CTG GAC CCC TCT CGA CTC GAG
AGT TCC GCT CCA CCG TTG GAC TTG CTC CGC TGT CGG CAT CCA GAA AT

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 30986.2

μg per OD: 31.7

Millimolar Extinction Coefficient: 976.3

nmoles per OD: 1.0

Purity	Desalt	OD's	89.3
Tm (1 M Na+)	100	$\mu\text{g's}$	2833
Tm (50 mM Na+)	78	nmoles	91
% GC	61	Coupling Eff.	98%
Notes:		90 μg \Rightarrow 100 μg	

Primer 6:

Primer Name: UBI HSPB VER.4B

Primer Number: D0373B12 (B12)

Researcher:

Primer Length: 97

Sequence (5' to 3'): P-T TTC TGG ATG CCG ACA GCG GAG CAA GTC CAA CGG TGG AGC GGA ACT CTC
GAG TCG AGA GGG GTC CAG AGG CAG CGA CAG AGA TGC CGT GCC GTC TGC

Molecular Weight $\mu\text{g}/\mu\text{mole}$: 31781.4

μg per OD: 29.6

Millimolar Extinction Coefficient: 1070.6

nmoles per OD: 0.9

Purity	Desalt	OD's	97.1
Tm (1 M Na+)	100	$\mu\text{g's}$	2883
Tm (50 mM Na+)	78	nmoles	90
% GC	62	Coupling Eff.	98%
Notes:		900 μg \Rightarrow 100 μg	

* - See Note about Quantities in Supporting Information.



LIFE TECHNOLOGIES.

Do Nucleobond preps of 5596, 5597,
4216, 4217, 4218 and 4219

Digest 5596 and 5597 w/ EcoRI as a check
make sure smaller frag. is ~2kb.

Digest 4216, 4217, 4218, 4219 w/ BglI / Salt
to check that the 168bp frag. is generated.


Digest 4216 w/ BcgI / XcmI to use as
accepting vector for ubiquitin versions 1-4.
Gel isolate on 1% agarose

Digest 5596 w/ NheI / NotI to isolate insert
(~1.6 kb) L4:BAASS. NotI and NheI
Gel isolate on agarose
cut in different buffer
Cut w/ NotI 1st & 2nd
and EtOH ppt.
Cut w/ NheI

Mini-preps on BAASS:NA #5 5648 - 5665
Digest w/ EcoRI / PstI. Cut 118 also
Run on 10% acrylamide


→ Anneal Ubiquitin versions 1, 2, 3, and 4
align together.
Heat to 95°C for 5 min then stick on ice
Check them on a 10% acrylamide gel

Run, pre-cut 3770 NheI / NotI on an agarose
gel to check it out. Katherine



Check Ubi variation w/ a 113/Not digest (first 5 of each)

1. 1Kb ladder
- 2-6 Ubi 1 7593 - 7597
- 7-11 Ubi 2 7561 - 7565
- 12-16 Ubi 3 7579 - 7583
- 17-21 Ubi 4 7597 - 7601
- 22 ~~7062~~ NH/Not 7062 vector
- 23 6439 NH/Not frag
- 24 1Kb ladder




All look fine. Send
these for sequencing

Sequencing data shows 7368 to be correct for
Gus/PPR


Make Avidin/PPR using 7368 as start
Digest 7424 (Avidin/3720) and 7368 (Gus/PPR)
with Bam^{HI}/Not

- Gel:
1. 1Kb ladder
 - 2-5. 7368 (Gus/PPR) Bam/Not
 6. Skip
 - 7-10. 7424 (Avidin) Bam/Not



Isolate large vector frag from Gus/PPR and the smaller
insert band from Avidin

Katherine Baifera



GUS ASSAY


PURPOSE: TO QUANTITATE THE AMOUNT OF GUS IN CORN SEED

MATERIALS: REACTION PLATE - COSTAR EIA/RIA
READING PLATE - NUNC FLUORESCENCE POLYSEEP
 MU - 4 METHYLUMBELIFERONE (SIGMA M-1508)
 MUG - 4 METHYLUMBELIFERONE β GLUCURONIDE (SIGMA M-9130)
 MICROBALANCE
 FLUORESCENCE MICROPLATE READER

PROCEDURE: USE PROTOCOL FOUND ON PAGE # 57 OF THIS NOTEBOOK (#58).

RESULTS: DATA FOUND BELOW. (BASED ON 20-MIN READINGS.)

<u>SAMPLE #</u>	<u>% TSP</u>	<u>SAMPLE #</u>	<u>% TSP</u>
GSE 12020-4	0.088	GSE 05030-1	0.087
-5	ND	-2	0.54
GSD 01120-1	ND	-3	0.61
-2	ND	-4	0.16
-3	ND	-5	0.06
-4	ND	U 0808-1	0.001
-5	ND	-2	0.002
GSE 15070-4	0.28	-3	0.007
U 05050-1	0.17	-4	ND
-2	0.015	-5	0.001
-3	0.010	U 07050-1	0.3
-4	0.174	-2	0.089
-5	0.010	-3	0.27
U 05090-1	0.043	-4	0.013
-2	0.014	-5	0.43
-3	0.001		
-4	0.001		
-5	0.004		
GSC 01010-1	0.006		
-2	0.010		
-3	0.009		
-4	0.60		
-5	0.48		



Investigator:

Book # 58

Chris Brook Date:

Witness:

Elizabeth Wilcox Date:

GUS Assay

SEE PURPOSE, MATERIALS, & PROCEDURE BELOW.

Purpose: To quantitate the amount of GUS in corn seed extracts.

Materials: Reaction Plate-Costar ELARIA, non-tissue culture treated 96-well flat bottom plate
 Reading Plate-Nunc Fluorescence Polystyrene 96-well black plate
 MU 4-methylumbelliferone (Sigma M-1508)
 MUG 4-methylumbelliferone 8-glucuronide (Sigma M-9130)
 Microbalance
 Fluorescence microplate reader (Molecular Devices Gemini)

Reagents: Lysis Buffer: 50 mM sodium phosphate pH 7.0, 1 mM EDTA, 10 mM BME

Note: 50 mM sodium phosphate is made by mixing 97 ml of Stock A (0.2M NaH_2PO_4 (27.6 g/L)) with 153 ml of Stock B (0.2M Na_2HPO_4 (53.6 g/L) and bringing to a final volume of 1.0 L with dH_2O .

Also note that the 10 mM BME should be added to an aliquot of the lysis buffer fresh daily, enough for that day's experiment.

Stop Buffer: 0.2 M Na_2CO_3 (21.2 g/L)

1 mM MU Standard Stock: 4.96 mg MU in 25 ml dH_2O (made fresh daily)
 20 mM MUG Substrate Stock: 7 mg MUG in 1.0 ml 95% ethanol (made fresh daily).

Procedure: Corn seed extracts should already be prepared and analyzed for total protein according to standard procedures.

In a reaction plate, equilibrate up to 10- μg of total protein in a total volume of 100 μl lysis buffer. Generally samples can be analyzed with 1 μg total protein. Samples should be analyzed in triplicate.

Add standard curve to triplicate wells diluted as follows:

10 μl of 1 mM MU standard stock is diluted with 90 μl lysis buffer.
 10 μl of this 1:10 dilution is further diluted with 90 μl lysis buffer to give a 1:100 dilution.

0 mM MU standard 100 μl lysis buffer / well
 1000 nM MU standard 12.5 μl of the 1:100 dilution + 87.5 μl lysis buffer / well
 10,000 nM MU standard 12.5 μl of the 1:10 dilution + 87.5 μl lysis buffer / well
 100,000 nM MU standard 12.5 μl of the 1 mM MU stock + 87.5 μl lysis buffer / well

Prepare the reading plates by pipetting 175 μl of Stop buffer into all wells of the plate. You will need a separate plate for each time point required. Generally we take readings at 0, 15, 30 and 60 minutes.

Dilute the 20 mM MUG substrate stock to 5 mM with lysis buffer. Add 25 μl of 5 mM MUG to every well including both standard and sample wells and mix to start the reaction. Immediately after adding the MUG, pipette 25 μl of solution from the reaction plate into a prepared reading plate. Place the reaction plate at 37 $^{\circ}\text{C}$ until the next time point. At each subsequent time point, pipette 25 μl of solution from the reaction plate into a prepared reading plate.

Reaction is stable for several hours once it has been stopped. Note that stopping the reaction is essential for fluorescence formation.

Plates are read at 360 nm excitation wavelength and 460 nm emission wavelength.

The unknown samples are read against the standard curve in nM MU and the amount of GUS in the samples is calculated as follows:

Average nM MU for each sample (Mean Value Column) / minutes reaction proceeded = nM MU / min * 60 min / hr = nM MU / hr. Note that if there is > 100 FU in the 0 time point reading of the average nM MU, that value must be subtracted from the average nM MU at each subsequent reading. This value is then corrected for the amount of protein added in the sample by dividing by the total protein added to give nM MU / hr / μg . This value is converted to %TSP by multiplying by 1.68×10^4 which is a conversion factor determined while at Pioneer.

A Quality Control sample (a known amount of GUS spiked into control corn seed extract) may be run on each assay to determine reproducibility of quantitation.



RESULTS: DATA FOUND BELOW. (10-MIN READINGS)

Sample#	%TSP	Sample#	%TSP	Sample#	%TSP
GSG 01040-1	0	GSG 01110-1	0.6	GSC 01060-1	0
-2	0.4	-2	0.04	-2	0
-3	0.6	-3	0	-3	0
-4	0.5	-4	0.04	-4	0
-5	0.4	-5	0.04	-5	4.8
GSD 02130-1	0.1	GSC 01070-1	0.4	GSC 01130-1	8.4
-2	0.7	-2	0.3	-2	0.1
-3	0.9	-3	0.3	-3	8.6
-4	0	-4	0.5	-4	5.0
-5	0.8	-5	0.001	-5	0.07
GSG 01020-1	0	GSC 01040-1	0.1	GSC 01110-1	0
-2	0	-2	0.5	-2	9.2
-3	0.12	-3	0.03	-3	0
-4	0	-4	0.03	-4	0
-5	0.2	-5	0.004	-5	9.6
GSC 01030-1	0				
-2	4.0				
-3	4.2				
-4	0.5				
-5	9.5				

Investigator:

Book # 67

Investigator: Chris Brooks Date: _____
Elizabeth Wilfong Date: _____

Witness: _____
 Witness: _____ Date: _____